

BoviBrom®: Efficacy in Reducing *E. coli* and *Salmonella* on Meat

KEY FINDINGS

- BoviBrom® at 270 ppm is as effective as hot water in a commercial carcass wash system
- BoviBrom did not cause “graying” of carcass lean tissue
- BoviBrom is effective using room temperature water
- BoviBrom is applied at near neutral pH, making it equipment- and worker-friendly
- BoviBrom can be reused in downstream applications, resulting in cost and water savings

OBJECTIVE

Evaluate the efficacy of BoviBrom as an antimicrobial intervention in reducing aerobic plate counts (APC), *Enterobacteriaceae* (EBC), *E. coli* O157:H7 and *Salmonella* on inoculated fresh meat surfaces.

STUDY DESIGN


Treatments were evaluated using two different systems: a commercial carcass wash cabinet and a model carcass washer. Sections of beef flank muscle and heart were inoculated with bovine fecal solution containing approximately 6_{Log} CFU/cm² of *E. coli* O157:H7 and *Salmonella*.

- Two treatments
 - Hot water – 185° F at nozzles, 20 psi, 12 sec (flank) and 28 sec (heart), 1.8 gal/min
 - BoviBrom – 77° F, 35 psi, 12 sec (flank) and 28 sec (heart), 2.6 gal/min
- Simulated for 300 head/hour chain speed
- BoviBrom pH level
 - 75 ppm – 6.84 pH
 - 175 ppm – 6.76 pH
 - 270 ppm – 6.71 pH

- Excised samples taken before and after treatment
 - After treatment, samples were measured immediately and again after being chilled for 48 hours at 39° F
 - Enumerated for:
 - APC
 - EBC
 - *E. coli* O157:H7
 - *Salmonella*

RESULTS¹

BoviBrom treatments reduced APC, EBC, *E. coli* O157:H7 and *Salmonella* by the same or slightly lower amounts relative to hot water treatment. After 48 hours of storage, the reduction of organisms by BoviBrom and hot water treatments did not change. The study demonstrated that BoviBrom spray washing is as effective as an antimicrobial intervention for beef carcasses and variety meats.



RESULTS (continued)

Commercial Carcass Wash Log Reduction of APC, EBC and *E. coli* O157:H7

	Treatment	APC	EBC	<i>E. coli</i> O157:H7
Flank	Hot Water	3.5*	3.9	1.8
	BoviBrom	3.3	3.1	1.8
Heart	Hot Water	4.1	4.1	2.2
	BoviBrom	3.6	3.4	2.1

*Means in the same column within tissue type did not differ significantly $P < 0.05$

Model Carcass Wash Log Reduction of APC, EBC, *E. coli* O157:H7, *Salmonella*

	Treatment	APC	EBC	<i>E. coli</i> O157:H7	<i>Salmonella</i>
Flank	Hot Water	3.0 ^d	3.3 ^d	2.3 ^d	2.5 ^d
	BoviBrom	2.5 ^e	2.1 ^e	1.5 ^e	1.3 ^e
Heart	Hot Water	3.8 ^x	3.4 ^x	2.4 ^x	2.8 ^x
	BoviBrom	3.0 ^y	2.7 ^y	1.9 ^y	2.3 ^y

^d and ^e, and ^x and ^y denote means in the same column within tissue type bearing the same letter do not differ significantly $P < 0.05$

For questions or to learn more about the Passport Food Safety Solutions product portfolio, call **515-334-8035** or visit **www.passportfoodsafety.com**.

The label contains complete use information, including cautions and warnings. Always read, understand and follow the label and use directions.

The sales and delivery of Passport Food Safety Solutions food-safety products will be expressly governed by and subject to the limited product warranty and other terms and conditions of sale available at www.PassportFoodSafety.com.

¹Kalchayanad N., et al.; 2009. Effectiveness of 1,3-Dibromo-5,5 Dimethylhydantoin on Reduction of *Escherichia coli* O157:H7– and *Salmonella*-Inoculated Fresh Meat. *Journal of Food Protection*. 72:151–156.

Passport® and the BoviBrom Application System™ are trademarks owned by Passport Food Safety Solutions, Inc. BoviBrom is a trademark of Albemarle Corporation.
 © 2016 Passport Food Safety Solutions, Inc.
 pasprt 2770-3

